



Granite Peaks Endoscopy 10150 S Petunia Dr (Building 2) Sandy UT 84092 801-619-1115

# 2-Day MiraLAX<sup>®</sup> Bowel Prep Instructions

- PURCHASE: MiraLAX or generic polyethylene glycol (21 servings) to mix with drink below
- Gatorade<sup>®</sup> 96 oz (NOTHING RED OR PURPLE) Sugar-free drinks (G2<sup>®</sup>, Pedialyte, Powerade Zero<sup>®</sup>, Propel<sup>®</sup> or water) are also options.

### 7 Days BEFORE Your Procedure:

• PLEASE FOLLOW INSTRUCTIONS FOR DIABETIC/WEIGHT LOSS GLP-1 MEDICATIONS AS NOTED ON THE BACK OF THIS SHEET.

## Two (2) Days BEFORE Your Procedure:

- **Beginning with a clear liquid breakfast**, drink at least 8 glasses (64 oz) of CLEAR liquids throughout the day and evening. Clear liquids include: Water, sports drinks, black coffee, tea, any soda (light or dark), broth, gelatin, popsicles, and juice without pulp. **NOTHING RED OR PURPLE.**
- **STOP all solid foods.** NO SOLID FOOD, no dairy, until after your exam.
- **STOP fiber supplements**, such as Metamucil.

# The Day BEFORE Your Procedure:

#### NOTICE: OUR OFFICE MAY CALL THE DAY PRIOR TO ADJUST YOUR ARRIVAL TIME AS WE FINALIZE THE CENTER'S SCHEDULE.

- Mix 7 servings of MiraLAX in 32 oz of drink. Repeat this for two more batches, and refrigerate.
- At 9:00am, drink one-third (32 oz) of the total MiraLAX/Gatorade solution over the next 2 hours.
- Continue drinking clear liquids throughout the day and evening.
- At 5:30pm drink the next third (32) oz of the total MiraLAX Gatorade solution over the next 2 hours.
- A soothing ointment (Desitin<sup>®</sup>, Vaseline<sup>®</sup>) applied to the anal area may prevent discomfort and skin breakdown.

## **Procedure Day:**

- **4 hours prior to your appointment,** drink the remaining prep over the next 2 hours.
- You may continue drinking clear liquids <u>until</u> 2 hours prior to arrival. **NOTHING BY MOUTH** for 2 hours before your appointment.
- Bowel movements should be watery and clear yellow or clear green in color, and you should be able to see the very bottom of the toilet bowl. If you are not passing clear stool or it is still brown 2 hours prior to arrival time, please call our nurses at 801-619-1115 for instructions.

THIS IS A SEDATED PROCEDURE. A RESPONSIBLE DRIVER MUST REMAIN IN THE FACILITY THROUGHOUT YOUR APPOINTMENT. (90-120 minutes total). NO EXCEPTIONS.

Find Additional Information and Instructions on Our Website

## www.GranitePeaksGl.com

©2023 Granite Peaks GI, LLC. All Rights Reserved. All other trademarks are property of their respective owners.





# **IMPORTANT INFORMATION** *Please call us with any questions at 801-619-1115*

### **MEDICATIONS**

 Take your regularly prescribed medications at least two (2) hours before you arrive. (Except blood thinners or diabetic/weight loss medication as described here).

### **DIABETICS-INSULIN**

- Test your blood sugar before coming in for your procedure and bring the reading with you.
- If you take oral diabetes medication or insulin, **DO NOT** take it the morning of your procedure.
- Please contact the prescribing physician for diabetic care if you have any questions regarding your oral medication or insulin dosages.

### **DIABETICS/WEIGHT LOSS PATIENTS - Semaglutide, GLP-1 Receptor Agonists**

#### 7 Days BEFORE Your Procedure:

- PATIENTS CURRENTLY TAKING A GLP-1 RECEPTOR AGONIST FOR <u>DIABETES OR WEIGHT LOSS</u> such as Ozempic<sup>®</sup>, Semaglutide, Mounjaro<sup>®</sup>, Wegovy<sup>®</sup>, Rybelsus<sup>®</sup>, Trulicity<sup>®</sup>, Victoza<sup>®</sup> or similar compounded or generic medication, should contact the prescribing physician for instructions to stop the medication at least 7 days prior to your procedure. For your safety, your procedure will be rescheduled if these medications are taken within 7 days of your scheduled appointment.
- Monitor your blood sugar levels while off the medication. Adjust insulin accordingly.

### **BLOOD THINNERS**

- If you are taking blood thinners, call your prescribing physician and ask if you may stop taking the medication for the required amount of time prior to your procedure.
- You may continue taking aspirin, ibuprofen and other anti-inflammatory medications.

### **BRING WITH YOU**

- Your inhaler, if you use one
- Your insulin, if you use it
- Photo ID and Insurance Card
- Leave valuables and phone/devices at home or with your driver
- List of all medications, vitamins and supplements you are taking (include dosage and last date and time each medication was taken before your arrival)
- Wear a short-sleeved shirt